

A LIST OF THE COLEOPTERA TAKEN IN ALASKA
AND ADJACENT PARTS OF THE YUKON
TERRITORY IN THE SUMMER OF 1924

BY H. C. FALL

Tyngsboro, Massachusetts

It was the writer's privilege to be one of a party of three to make a trip to this wonderland of the North during the summer of 1924. While the excursion was primarily one of sight-seeing and general recreation, the writer, from long habit, kept always an eye open for possible coleopterous windfalls, and as opportunity offered, a few minutes, a few hours, or even the better portion of a day in places where a long enough stop permitted, were devoted to special search for beetles. The results in the aggregate seem worthy of record, in view of the general interest taken in collections from this somewhat remote region.

In 1857 Le Conte remarked that at the close of the activities of the Russian entomologists, the fauna of the region explored by them was more completely developed than that of any other part of the continent. This, of course, is no longer true, even if we confine ourselves to the coast region in which most of the early collecting was done; while if the entire territory of Alaska be considered, it would perhaps be nearer the truth to say that its fauna is less completely known than that of any other equal area of our country.

The total number of species of Coleoptera listed from Russian America at the time of Mannerheim's last supplement in 1853 was 540. Hamilton's Alaskan list with supplement in 1894 enumerates 578 species, and the few scattering additions since that time can hardly bring the total much, if any, above 600 species. It is quite probable that there are now in collections, notably that of the National Museum, a considerable number of additional species, but these, even if worked up, have not yet been recorded.

On the present trip, as nearly as can now be determined, 448 species were taken within, or adjacent to, the Alaskan boundary, of which 117, exclusive of unidentified *Stenus* and *Aleocharini*, appear to be new to the Alaskan list. Of these, 61 occurred in the coast region and 64 in the Yukon Valley,

8 species being common to both regions. The fact that nearly half of these 117 species are from the well-explored coast region is rather surprising, since it was to be expected that the great majority of future additions to the Alaskan fauna would be from the interior valleys and mountains, which, entomologically speaking, are relatively unworked. The disparity in this case would indeed have been much greater if the numerous species taken at Dawson, Y. T., where considerable collecting was done, had been included, for it is practically certain that a very large proportion of the species occurring at this point must also occur across the international boundary only fifty miles distant, where the topographical and climatic conditions along the Yukon are virtually the same.

In the interest of economy of space, I shall not attempt here to speak of collecting conditions and experiences, or, except for the table of dates below, to give the itinerary of the trip, further than to say that we left Tyngsboro, Massachusetts, on June 5, via Montreal and the Canadian Pacific, arriving at Skagway June 18, after making the incomparable trip up the "Inside Passage" from Vancouver. On August 2 we boarded the steamer at Ketchikan for Prince Rupert, the return journey across the continent being made thence via the Canadian National Railway to Montreal, arriving at Tyngsboro August 10.

The principal dates on which collecting was done are given once for all in the subjoined table:

Wrangel, Alaska, June 17.	Anchorage, Alaska, July 10-14.
Skagway, Alaska, June 18-20.	Seward, Alaska, July 16-20.
Skagway, Alaska, July 27.	San Juan, Evans Island,
White Horse, Y. T., June 21-22.	July 20.
Dawson, Y. T., June 24-30.	Childs Glacier, Alaska,
Eagle, Alaska, July 1.	July 22.
Fort Yukon, Alaska, July 2.	Port Althorp, Alaska, July 23.
Nenana, Alaska, July 6.	Juneau, Alaska, July 25-26.
Fairbanks, Alaska, July 7.	Sitka, Alaska, July 28.
Mount McKinley Park, July 8-9.	Scow Bay, Alaska, July 30.
	Kasaan, Alaska, July 31.
	Ketchikan, Alaska, August 1.

Throughout the 1400-mile trip down the Yukon and up the Tanana Rivers daily stops were made for fuel at wood-piles

along the banks, at which nearly always some beetles were taken.

Species new to the Alaskan list are marked with an asterisk (*).

List of Species

FAMILY CICINDELIDÆ

Cicindela longilabris Say. White Horse, Y. T. Several seen, but having no suitable net no effort was made to secure specimens. No tiger beetles whatever were observed in Alaskan territory.

FAMILY CARABIDÆ

Trachypachus inermis Mots. Skagway; White Horse.

Cychnus (Pemphus) angusticollis Mann. Seward, Sitka.

Cychnus (Brennus) marginatus Fisch. Juneau; Skagway; Kasaan; Ketchikan. Common everywhere near the coast; individuals from the last two localities are considerably larger than those from farther north.

Carabus tædatus Fab. Skagway.

Elaphrus riparius L. Skagway; McKinley Park; Fort Yukon, June 27 (Kusche).

Elaphrus pallipes Horn. Skagway; Dawson. All typical in coloration.

Loricera 10-punctata Esch. Scow Bay. A single example.

**Opisthius richardsoni* Kby. Skagway; Dawson; Fort Yukon, June 27 (Kusche). Frequent along the river at Skagway, varying greatly in color, coppery, green, or nearly black. There are specimens of this species in my collection from Dolomi, Alaska, collector unknown to me.

Notiophilus borealis Harris. McKinley Park.

**Notiophilus simulator* Fall. Anchorage. A single example; also taken by me at Jasper Park, Alberta, on the return trip.

Nebria metallica Fisch. Skagway; Seward; Child's Glacier; Sitka; Juneau.

Nebria sahlbergi Fisch. Seward; Juneau. Two forms referable to this species were taken on the trip; one of narrow thorax and somewhat more elongate form at Juneau and Seward is probably the true *sahlbergi*. Of this a variety with violaceous elytra (probably *violaceus* Mots.) and an-

other with pale legs occurred. The other form is represented by a single example taken at Goldstream, Vancouver Island; it is a little stouter with broader thorax, agreeing closely with specimens in my collection from Washington State and the Rocky Mountains of New Mexico. This form, in its broader thorax, approaches *mæsta* Lec., which, I think, can hardly be a varietal form of *sahlbergi*, although so recorded in the books.

Nebria bifaria Mann. A single typical specimen with bicolored legs was taken at McKinley Park.

Nebria gebleri Dej. Juneau.

Nebria viridis Horn. Dawson; Yukon River (Sheep Creek below Eagle); Tanana River.

Nebria hudsonica Lec. White Horse. Two examples.

Nebria testaceipes Cys. Dawson. One example.

**Dyschirius secretus* Fall, n. sp.

Slightly smaller than, but of the general form and aspect of, *integer* Lec., near which it must be placed.

Body black, bronzed; legs and antennæ black. Head four-fifths as wide as the prothorax; front feebly transversely impressed; epistoma broadly and deeply emarginate. Prothorax subglobose, slightly wider than long, sides broadly, evenly arcuate. Elytra oblong-oval, three-fifths longer than wide; striæ entire, rather deeply impressed and strongly punctured before the middle, becoming much finer and more finely punctured on the declivity; eighth stria finely impressed, distinctly punctate throughout; base not margined; third interspace tripunctate, a single apical setigerous puncture. Basal joint of hind tarsi twice as long as wide and evidently shorter than the two following combined. Length, 2.5 to 2.75 mm.; width, .8 to .85 mm.

Anchorage, Alaska, seven examples, all taken in damp moss on low ground, July 13-14, 1924.

In *integer*, with which the present species is compared, the size is slightly larger, the legs and antennæ are never really black, but rather dark rufous to rufopiceous, the basal joint of the hind tarsus is more elongate, at least three times as long as wide and fully as long as the two following; apex of elytra with two setigerous punctures. There is a strong probability that *integer* is distinct from the European *æneus* with which it has been united, judging from two examples of the latter sent me by Mr. Edmund Reitter. In *nigriceps*, with which the present species might also be compared, the size is larger, the

elytra more finely striate and punctate, the eighth stria obsolete, and the elytral apex apparently with two setigerous punctures; the basal joint of the hind tarsus is also more elongate, as in *integer*.

Only two species of *Dyschirius* have hitherto been recorded from Alaska, viz., the *frigidus* and *transmarinus* of Mannerheim. The former was described from the Kenai Peninsula, and my Anchorage species might well be expected to be the same. I find, however, that my specimens do not fit the description very well, the size of *frigidus* (3.5 mm.) being materially larger, while the elytra are said to be scarcely wider than the thorax and the elytral striæ evanescent shortly behind the middle. If *frigidus* really equals *æneus*, as it stands in our lists, it certainly is different from the Anchorage species. *Transmarinus* was described from a single example from Sitka; it is even larger than *frigidus*.

Asaphidion yukonense Wickh. White Horse; Dawson.

Bembidion simplex Haywd. McKinley Park.

Bembidion planiusculum Mann. Sitka.

Bembidion complanulum Mann. Dawson; McKinley Park; Seward; Sitka; Kasaan.

Bembidion funereum Lec. Skagway; Dawson.

Bembidion mæklini Haywd.? Sitka.

Bembidion quadrifoveolatum Mann. Kasaan.

***Bembidion yukonum* Fall, n. sp.**

Form and size about as in *grapi*, but rather larger and a little stouter. Black, distinctly green-bronzed, antennæ at base and legs rufous; surface above strongly shining throughout, without alutaceous sculpture except very feebly at elytral apex. Antennæ scarcely attaining the middle of the length, intermediate joints about two and a half times as long as wide. Head a little wider than the thoracic apex; eyes moderate. Prothorax subcordiform, one-fifth wider than long, base slightly narrower than apex; sides strongly evenly rounded from apex to basal constriction, which is parallel-sided for a distance about equal to one-fifth the basal width; base angles sharply rectangular, evidently but not very strongly carinate; basal impressions deep, with a few rather coarse punctures. Elytra one-half wider and three times as long as the thorax, one-half longer than wide; sides broadly arcuate, humeri well marked, not angulate; striæ feebly impressed near the suture; stria punctures moderate, becoming finer posteriorly, very fine but traceable near the apex; those of seventh stria very fine in apical half; intervals flat, the third with two setiger-

ous punctures adjacent to the third stria at approximately the basal and apical thirds. Length, 4.75 mm.; width, 1.9 mm.

Described from a single male specimen taken by the writer at Dawson, Yukon Territory, June 30, 1924.

This species belongs to the *ustulatum* group, and is apparently nearest *insopitum* Csy. The latter is unknown to me, but according to description is a considerably larger species with more transverse thorax, deeply striate elytra, the seventh stria less distinctly punctured basally. *Grapi* is a rather smaller and narrower species, with much finer elytral sculpture, the legs typically black, though in two Greenland examples in the Le Conte collection and in many North American examples so referred, they are more or less distinctly rufous.

Bembidion bimaculatum Kby. Skagway; Dawson; Nenana.

There occurred at Nenana, together with the typical form, what I take to be a variety of this species with smaller thorax, browner elytra with or without apical spot, and finer striæ.

****Bembidion mckinleyi* Fall, n. sp.**

Black, above with feeble deep-green surface luster, beneath black, femora piceous; trochanters, tibiæ, and tarsi reddish-brown. Upper surface polished throughout, with faint traces of alutaceous sculpture only at elytral apex. Antennæ black, basal joint rufescent, about half as long as the body, slender, intermediate joints not quite three times as long as wide. Head three-fourths as wide as thorax, eyes not at all prominent. Prothorax nearly one-third wider than long, sides rounded anteriorly, sinuate posteriorly, becoming nearly parallel in basal fifth or sixth; hind angles rectangular, side margins very finely reflexed; disc convex, median line well impressed and nearly entire; foveæ only moderately deep, flat or slightly tumid at bottom, with a few obsolete punctures, hind angles not distinctly carinate. Elytra fully three and one-half times as long and three-fourths wider than thorax, three-fifths longer than wide, humeri rather wide, narrowly rounded, sides diverging slightly and nearly straight in basal two-fifths, greatest width at or very slightly behind the middle; striæ finely impressed, moderately strongly and closely punctured basally, more feebly so posteriorly, nearly disappearing at apex; seventh stria represented by a few fine punctures near the base. Length, 5.5 mm.; width, 2.25 mm.

Described from a single pair taken in McKinley Park, Alaska, July 8-12, 1924, by my fellow-traveler, Mr. C. A. Richmond.

This species belongs to the *ustulatum* group, and by its narrow thorax is somewhat remindful of *scopulinum*. It is, how-

ever, much larger and without maculation. Careful measurement shows that the prothorax is not quite as wide at apex as at base. On this basis it would, by Hayward's table, be associated with *nevadense* and *consanguineum*, to neither of which is it at all similar. By Casey's table it would fall near *humboldtense*, which is even more remotely related. If the thorax were regarded as narrower at base than at apex (this relation is at best a rather elusive one) it would fall between *scopulinum* and *canadense* in Hayward's table, which is perhaps the most satisfactory position for it.

Bembidion rupicola Kby. (*lucidum* Lec.). Skagway; Yukon and Tanana Rivers. *Rupicola* has long stood on our lists as a synonym of *ustulatum*, but wrongly so. The synonymy here announced is made certain by an examination of one of Kirby's cotypes kindly sent me by Mr. Arrow.

Bembidion scopulinum Kby. Anchorage; Dawson.

Bembidion picipes Kby. (*grapi* Auct.). Skagway; Dawson; Yukon River (Sheep Creek, below Eagle); Tanana River; McKinley Park. This is the species we are accustomed to call *grapi*, and probably correctly so, although I have not been able to compare with European specimens. Except for darker legs, a Sheep Creek specimen is almost an exact counterpart of one of the cotypes of Kirby's *picipes*, which I have had the privilege of examining. There can be no doubt that the other examples taken are the same thing, and that the species long standing as *picipes* on our lists, following Le Conte's misinterpretation, must henceforth be known as *plagiatum* Zimm.

Bembidion dentellum Thumb. (*incrematum* Lec., *arcuatum* Lec.). Dawson.

**Bembidion coloradense* Haywd. Anchorage.

Bembidion nigripes Kby. Skagway; Dawson.

Bembidion intermedium Kby. I did not take this species, but there are examples in my cabinet collected by Kusche at Fort Yukon, June 27, 1916. This is the true *intermedium* of Kirby, which seems never to have been correctly interpreted by our systematists. It is very closely allied to *nigripes* (perhaps not distinct), and is similarly brightly variegated with yellow. The much less ornate *intermedium*

of the Hayward monograph is an entirely different thing, for which Le Conte's name *rapidum* must be restored, unless on investigation some still older name comes to light.

Bembidion convexulum Haywd. Dawson. Two examples.

Bembidion atrolucens Csy. White Horse.

Bembidion versicolor Lec. Wrangel.

Bembidion dubitans Lec. Dawson.

Bembidion mutatum G. & H. Dawson.

**Bembidion sulcatum* Lec. Anchorage. A single example.

****Bembidion concurrens*** Fall, n. sp.

Body oblong, suboval, rather strongly convex; black, polished throughout, elytra with a small nubilous external pale spot at apical fourth, the apex barely visibly pallescent. Antennæ black, basal joint rufescent; median joints twice as long as wide. Head only slightly narrower than thorax; eyes rather large and prominent. Prothorax one-half wider than long, apex evidently wider than base, sides broadly arcuate in front, posteriorly sinuate and subparallel in about the basal sixth, the angles sharp and just visibly more than right; basal foveæ deep, not evidently punctured, bordered externally by a fine acute carina. Elytra not quite one-half longer than wide, five-eighths wider than thorax, parallel, sides feebly arcuate in basal half; striæ finely impressed internally, scarcely at all so in outer half; stria punctures moderate, becoming much finer but still quite distinct on the declivity and laterally in apical half. Legs very nearly black. Length, 3.25 mm.; width, 1.35 mm.

Alaska. Scow Bay (north of Wrangel) two males; Skagway, one female. The type is a male from Scow Bay.

This species belongs to the *assimile* group and in form most nearly resembles *anguliferum*, though materially smaller and with elytra much less finely punctate-striate. The species of this group form a difficult study, and are considerably more numerous than indicated by Hayward's very conservative treatment. The present species by a rigorous interpretation of Casey's table would fall near *invidiosum*, in which, as in *cautum*, the elytra are not truly black (as they are in the present species) and the legs are rufous. The elytra are also relatively shorter and more coarsely striate than in *concurrens*. It is possible that *invidiosum* is not really distinct from *cautum*, and I feel tolerably sure that *digressum* Csy. is the same as *connivens* Lec.

Bembidion sp. dub. (*assimile* group). Dawson; Seward.

- Bembidion oblongulum* Mann. Seward.
Patrobis septentrionis Dej. Skagway.
Patrobis (Platidius) aterrimus Dej. Ketchikan.
Trechus chalybæus Dej. Skagway; Anchorage; Seward;
 Kasaan.
Pterostichus crenicollis Lec. Ketchikan.
Pterostichus amethystinus Dej. Kasaan.
Pterostichus validus Dej. Juneau; Kasaan; Ketchikan.
Pterostichus castaneus Dej. Scow Bay; Kasaan; Ketchikan.
Pterostichus adstrictus Esch. Skagway; Juneau; Anchorage;
 Seward; Scow Bay; Kasaan; Dawson. Nearly all the names
 under *Bothriopterus* of the Leng list apply to more or less
 trifling variations of this ubiquitous northern species.
Pterostichus (Cryobius) riparius Dej. Skagway; Seward;
 Juneau.
Pterostichus (Cryobius) hudsonicus Lec. Childs Glacier.
Pterostichus (Cryobius) brevicornis Kby. McKinley Park;
 Dawson.
Pterostichus (Cryobius) sp. near *mandibularis*, and perhaps not
 different. McKinley Park; Tanana River.
Pterostichus (Cryobius) sp. McKinley Park; Eagle; Dawson.
 I am unable to place this, even tentatively.
Pterostichus (Cryobius) sp. Anchorage. A single specimen
 allied to the preceding, but with entirely black palpi, an-
 tennæ, and legs.
Amara (Cyrtonotus) eschscholtzi Chd. Dawson.

****Amara (Cyrtonotus) consueta* Fall, n. sp.**

Oblong, moderately elongate, form nearly as in *eschscholtzi*. Black, head and thorax polished; elytra minutely alutaceous, moderately shining in the male, slightly duller in the female; antennæ and legs ferruginous. Head not quite two-thirds as wide as prothorax. Prothorax not quite one-half wider than long, base very slightly wider than apex; sides broadly, evenly rounded anteriorly, briefly sinuate before the nearly rectangular hind angles; disc moderately convex, anterior transverse impressed line feeble; median line distinct, abbreviated at both ends; disc in front completely devoid of punctures, basal impressions deep, a little oblique, numerous punctate; hind angles obtusely carinate; base at middle impunctate, side margins narrowly, finely punctured. Elytra (♂) about one-fifth wider and not quite three times as long as the thorax and about three-fifths longer than wide; finely striate, striæ finely but distinctly punctate, absolutely so near apex. Body beneath black, meso- and metepisterna and

sides of the metasternum and basal segments of abdomen with a few fine punctures. Middle and hind thighs bisetose on inner margin; tarsi with first three joints feebly grooved at sides. Length, 10.5 to 11 mm.; width 3.9 to 4 mm.

Described from six examples taken by the writer at Skagway, Alaska, June 18-21, 1924.

This species is, by a strict interpretation of Hayward's table, to be associated with *infausta* Lec., and is undoubtedly closely allied with both that species and *eschsoltzi*, from both of which it seems constantly separable by the total lack of anterior discal punctuation of the thorax. It seems highly probable that we have too many described species already in this genus, nevertheless I am unable to attach any of the published names to the present species.

Amara littoralis Esch. Skagway; Ketchikan.

Amara remotestriata Dej. White Horse.

Amara interstitialis Dej. Dawson.

**Amara polita* Lec. Eagle.

Calathus ingratus Dej. Skagway; Nenana; Anchorage.

Calathus advena Lec. Childs Glacier.

Platynus stygicus Lec. (|| *maurus* Mots.). Anchorage.

Platynus perforatus Lec. Dawson.

Platynus bogemanni Gyll. Skagway; Dawson.

Platynus quadripunctatus DeG. Anchorage.

Platynus n. sp. ? near *sordens* Kby. Anchorage.

**Platynus nigriceps* Lec. Fairbanks.

Cymindis cribricollis Dej. White Horse.

Harpalus fulvilabris Mann. Eagle.

Harpalus pleuriticus Kby. Dawson.

**Harpalus laticeps* Lec.? Fairbanks. A single non-typical example, somewhat crushed, picked up from the sidewalk.

**Harpalus cordifer* Notm. Juneau; taken also at Goldstream, Vancouver Island.

Harpalus basillaris Kby. White Horse; Eagle.

**Harpalus pacificus* Fall, n. sp.

Body oblong, rather markedly convex, black, strongly shining throughout, the elytra very finely alutaceous. Antennæ and trophi ferruginous, the former scarcely passing the hind angles of the thorax, their intermediate joints fully twice as long as wide; head not quite three-fourths as wide as thorax; eyes moderately prominent. Prothorax slightly more than one-half wider than long; base

a little wider than apex; sides subevenly arcuate, a little more strongly so in front; basal angles obtuse, their apices somewhat blunted but not distinctly rounded; basal foveæ rather strongly, closely punctured, the adjoining surface externally sparsely punctured, the punctures nearly absent at middle of base; sides strongly convexly declivous anteriorly, narrowly and feebly deplanate posteriorly from the middle, the marginal bead translucent posteriorly. Elytra not quite one-half longer than wide, sides parallel; striæ fine, intervals flat with scattered, very minute punctures. Body beneath black, the epipleura not at all paler. Legs rather stout; femora and tips of tibiæ black; tibiæ except at apex and tarsi rufous. Length, 8.5 mm.; width, 3.5 mm.

Ketchikan, Alaska, a single male (type); Scow Bay, one female; Skagway, one male doubtfully referred.

This species in its typical form agrees nearly with the description of *fugitans* Csy. in many respects, including the color of the legs, but in that species the form is said to be unusually depressed, the hind angles of the thorax nearly right, and the pronotal foveæ very shallow and sparsely punctured, none of which statements are at all applicable to the present species.

The Scow Bay and Skagway specimens differ from the type in having the legs predominantly rufous, the tips of the femora and tibiæ only being lightly infuscated, the sides of the thorax are also less rounded. In the Scow Bay female the elytra are dull. All three specimens have the mentum emargination virtually without a tooth, although there is an extremely feeble, broadly rounded, prominence detectable on close scrutiny.

Harpalus rufimanus Lec. Dawson.

Trichocellus cognatus Gyll. Seward.

Tachycellus nigrinus Dej. Skagway; Juneau; Scow Bay; Wrangel.

FAMILY DYTISCIDÆ

Cœlambus borealis Fall, n. sp.

Elongate oval, widest at middle of the length, sides of thorax nearly continuous with those of the elytra. Head and thorax flavotestaceous, the former with a dark spot adjacent to each eye and the hind margin narrowly blackish, the latter with the anterior margin narrowly dark, and a small central spot which is diffusely produced to the hind margin. Elytra with the disc broadly fuscoluteous, due to the diffusion of the customary markings, which are, however, somewhat defined when the elytra are viewed very obliquely in the direction of their length. Body beneath black except the head, thorax, and epipleura; legs rufotestaceous; tarsi blackish. Head

alutaceous and sparsely punctulate. Prothorax and elytra polished throughout, the former finely and sparsely punctured on the disc, more closely toward all the margins, more strongly so post-medially; elytral punctures nearly equal in the posterior half where they are separated by about half their own diameters, in the anterior half they are obviously, though not very conspicuously, unequal and somewhat less closely placed. Prothorax one-half wider than the head, the sides moderately divergent and almost straight throughout. Elytra twice as wide as the head and one-half longer than wide. Sides of coxal plates and of the abdomen basally, coarsely closely punctate, the punctures separated by much less than their own diameters; surface of coxal plates strongly alutaceous.

Male. Four anterior tarsi strongly dilated, the second joint of the protarsus visibly wider than the first and as wide as the tibiae at apex; anterior protarsal claw a little thicker and just visibly shorter than the posterior and more strongly curved basally. Length, 3.7 mm.; width, 1.85 mm.

Dawson, Yukon Territory. A single male taken by the writer June 30, 1924.

This species is very closely allied to *patruelis*, and is possibly only a variety thereof. It differs from all specimens of a numerous series of *patruelis* in my collection by the almost perfectly straight sides of the prothorax and by the strongly alutaceous coxal plates, these being polished and without alutaceous sculpture, except in the dull females, in *patruelis*. The punctuation of the coxal plates in *patruelis* is also less dense than in the present species, the elytral disc is more uniformly punctured, and the tarsi only rarely show any tendency to infuscation.

Hydroporus planiusculus Fall. Scow Bay.

Hydroporus notabilis Lec. Anchorage. Specimens vary considerably in size and indicate the identity of *notabilis* and *arcticus*.

Hydroporus nigellus Mann. Anchorage; Seward; Skagway.

Hydroporus tartaricus Lec. McKinley Park.

Hydroporus melanocephalus Gyll. Dawson; Tanana River; Fairbanks.

Hydroporus polaris Fall. Dawson. A single male of this rare species.

**Hydroporus fuscipennis* Schaum. Fairbanks.

Hydroporus striola Gyll. Dawson.

***Hydroporus yukonensis* Fall, n. sp.**

Allied to *12-lineatus* and *semiclarus*. Moderately elongate, oval; upper surface pale testaceous with blackish markings as follows:

Head with a subtriangular spot each side adjacent to the eyes, these spots confluent with each other and with a posterior marginal line at the middle of the occiput; prothorax with a small transverse spot each side of the middle in the posterior transverse impression and another still smaller at the middle of the sublateral longitudinal impression; elytra each with six narrow discal vittæ and two or three exterior longitudinal dashes, the black vittæ a little wider than the pale stripes between them. Prothorax wider at base, twice as wide as long, and about one-half wider than the head; sides oblique and nearly straight from the base almost to the apical angles; side margins fine but distinct. Elytra about four and one-half times as long, and one-half wider than the thorax, and a little less than one-half longer than wide; sides markedly discontinuous with those of the thorax, subparallel and broadly arcuate in basal three-fifths; feebly sinuate before the apex in the male, obtusely angulate before an apical sinus in the female; the sutural angles acute and well defined in both sexes. Upper surface throughout micro-reticulate and sparsely, finely punctulate. Body beneath black, similarly micro-reticulate and sparsely punctulate; margins of hypomera and epipleura testaceous. Antennæ and legs testaceous, the apical parts of the terminal joints of the palpi and outer joints of the antennæ and the basal two-thirds of the femora infuscate.

In the male the protarsi are a little wider than in the female, with the second joint evidently wider than the first (scarcely wider than the first in the female), and the protarsal claws slightly less slender, though mutually equal and scarcely longer than in the female. The last ventral segment is not appreciably modified in either sex. Length (type male), 4.5 mm.; width, 2.3 mm.

Described from a single pair taken by the writer at Dawson, Yukon Territory, June 24, 1924.

This species is very nearly allied to *semiclarus*, and may prove to be only a smaller Northern race of that species. The two agree in having the male protarsal claws slightly thickened, but the present species, in addition to the smaller size, differs in its somewhat smaller thorax with nearly straight converging sides, the sides being much more rounded in *semiclarus*; *12-lineatus* differs from both by the protarsal claws of the male not at all thickened, the entirely pale femora and the more ample prothorax with sides less strongly discontinuous with those of the elytra.

***Hydroporus alaskanus** Fall, n. sp.

Similar to the preceding species in color, markings, general form and structural features, except as follows: Form a little more elongate, more than twice as long as wide (less than twice as long as wide in *yukonensis*). Head three-fourths as wide as the prothorax, the latter with sides straight and parallel in basal half to three-

fourths, extremely finely margined; hind angles rectangular; disc with two anterior transverse black dashes parallel with those in the posterior depression; elytra with sides even more strongly discontinuous with those of the thorax. In the male the protarsi are as in *yukonensis*, except that the tarsal claws are virtually as slender as in the female; the last ventral is broadly transversely impressed medially, the apical half flattened or feebly impressed longitudinally at middle. In the female the apex of the last ventral is produced in the form of a short broad deflexed process with converging sides and broadly truncate extremity. Length, 4.5 to 5.3 mm.; width, 1.9 to 2.4 mm. Ratios of length to width in four measured examples are 2.37 (δ); 2.2, 2.2, 2.1 (♀).

Described from one male and seven females, all taken by the writer at Skagway, Alaska, June 18-21, 1924. It is quite distinct from the preceding species by the above characters, and from all allied forms by the peculiar apical process of the last ventral in the female.

***Hydroporus recticollis** Fall, n. sp.

Closely allied to the preceding two species, but more especially to *alaskanus* with which it agrees in the short parallel-sided thorax with rectangular hind angles, the nearly straight sides extremely finely margined. The form is more strongly elongate, the thorax smaller than in either of the preceding two species and not much wider than the head; elytra from five and one-half to six times the thoracic length. Black markings of the upper surface heavier than in the preceding species, the thoracic disc becoming diffusely clouded, leaving the margins and a narrow central line pale; the black elytral vittæ much wider than the intervening yellow lines and the external fragmentary vittæ more developed. There is a tendency here, as in all species of the group, for the first and second elytral vittæ to become confluent in a small antemedian spot. The epipleura are entirely blackish. In the male, as in the preceding species, the protarsi are perceptibly more dilated than in the female, the second joint a little wider than either the first or third; the claws mutually equal, slender and not differing sexually. The last ventral is longitudinally medially impressed in apical half in the male. In the female the last ventral is rather abruptly flattened or impressed around the apical margin, which is at middle a little deflexed, but not distinctly produced. The subapical marginal angulation of the elytra in the female is a little more prominent, being nearly rectangular in the present species, obviously obtuse in the two preceding. Length (male type), 5.2 mm.; width, 2.3 mm. In two females the dimensions are 4.9 and 2.15 and 4.8 by 2.15 mm.

Described from one male and two female examples, taken by the writer at Seward (Kenai Peninsula), Alaska, July 18, 1924.

The five species now known from our fauna, in this restricted group having the sides of the elytra of the female subapically angulate, may be tabulated as follows:

Prothorax wider at base than at middle.

Protarsal claws not differing appreciably in the sexes, femora entirely pale.....*12-lineatus*

Protarsal claws slightly thickened in the male; femora blackish in basal half to three-fourths.

Sides of thorax arcuate throughout.....*semiclarus*

Sides of thorax nearly straight.....*yukonensis*

Prothorax not appreciably wider at base than at middle, the sides virtually straight and parallel in at least basal half; protarsal claws of male not at all thickened.

Last ventral of female with a broad deflexed apical process*alaskanus*

Last ventral of female without broad apical process; general form a little more elongate; prothorax smaller and with broad diffuse discal cloud.....*recticollis*

Agabus hypomels Mann. Ketchikan; Scow Bay; San Juan (Evans Island).

**Agabus inscriptus* Cr. Skagway. I have also in my cabinet a specimen taken at Homer by W. S. McAlpine of Detroit.

**Agabus* sp. Allied to *canadensis*; a single male taken at Fairbanks. It is possible that the *lecontei* Cr. of the Hamilton list may be this.

Agabus infuscatus Aubé. Anchorage.

Agabus anthracinus Mann. San Juan (Evans Island).

**Agabus nigroæneus* Er. Fairbanks.

****Agabus kenaiensis* Fall, n. sp.**

Obtusely oblong oval, black, antennæ and legs brownish, the thighs darker, especially the posterior. Body only moderately convex, surface rather strongly shining, minutely alutaceous and with a feebly impressed system of much larger reticulations which are quite irregular in form. Punctuation apparently lacking, except for the usual irregular finely setigerous elytral series and similar irregular series along the front margin and the exterior third of the rear margin of the prothorax. Closer inspection, however, reveals numerous scattered very minute punctures mostly at the intersections of the fine reticulating lines. Head with two faint rufous occipital spots. Prothorax a little less than three times as wide as long, sides feebly arcuate, nearly continuous with the sides of the elytra. Margin moderate, base just perceptibly sinuate each side of the middle. Elytra two-thirds as long as wide, slightly wider at middle than at base, sides subparallel in basal two-thirds. Body beneath alutaceous-reticulate, with a few fine punctures; ventral surface with some

widely spaced oblique very fine striæ, especially on the second segment. Prosternal process roof-shaped; hind coxal plates distant by rather less than half their own length from the middle coxal cavities; hind tibiæ with a few punctures along their inner margin.

Male. Anterior tarsi feebly dilated, somewhat compressed, clothed beneath with rather long squamules bearing at their tips a few small inconspicuous palettes; prosternal claws a little elongate, slender, equal, the posterior with a more pronounced basal sinuation of the inner margin. Length, 6.25 mm.; width, 3.3 mm.

Described from two completely similar male specimens taken by the writer at Anchorage, Alaska, July 14, 1924. They occurred in wet moss.

This species is nearly allied to *confertus*, differing in its much smaller size, somewhat narrower form, elytra without a trace of a sublateral pale spot, last ventral segment of male nearly destitute of the longitudinal strigæ so conspicuous in *confertus*. In the present species the posterior protarsal claw shows the more conspicuous basal sinuation, while in *confertus* it is the anterior claw.

**Agabus gelidus* Fall, n. sp.

Moderately elongate, oblong oval, barely perceptibly inflated posteriorly, elytra and thorax nearly continuous in outline; black, moderately shining; antennæ dark rufous; legs piceo-rufous. Very similar in most of its characters to *confertus*, with which it agrees nearly in size and form; the sculpture above consists as in *confertus*, of a system of rather large irregular reticulations within which the surface is more finely reticulate, with sparse, very fine scattered punctures, which may lie either within the meshes or on the reticulating lines. As in *confertus*, the metasternal groove for the reception of the tip of the prosternal process is quite rudimentary. The series of punctures along the inner margin of the hind tibiæ are very few in number, and the present species differs otherwise from *confertus* in the narrower side margins of the prothorax, and notably in the much narrower wings of the metasternum between the middle coxal cavities and the hind coxal plates, this distance being in the present species less than half the distance across the coxal plate, while in *confertus* it is obviously greater than half the width of the coxal plate. The sublateral yellow spot of the elytra is wanting in *gelidus*, and the longitudinal strigation of the last ventral in the male is only feebly developed, there being, however, a broad median impressed area extending from apex nearly to base of the segment, which seems to be wholly wanting in *confertus*. The protarsi of the male are similarly but noticeably more strongly dilated and compressed in *gelidus*, the protarsal claws elongate and sinuate on the inner margin in about the same degree. Length, 8 mm.; width, 4.25 mm.

Described from a single male specimen taken by the writer at Fairbanks, Alaska, July 7, 1924.

Agabus tristis Aubé. Skagway; Sitka; Wrangel; Ketchikan.

Ilybius quadrimaculatus Lec. Wrangel.

Rhantus divisus Aubé. Skagway.

FAMILY HYDROPHILIDÆ

Ochthebius holmbergi Mäkl. Anchorage.

Helophorus inquinatus Mann. Skagway.

**Helophorus lacustris* Lec. ?, McKinley Park.

**Helophorus nitidulus* Lec. Fairbanks.

Hydrobius fuscipes L. Fairbanks.

**Hydrobius scabrosus* Horn. Ketchikan.

**Paracymus moratus* Horn, *var.* Skagway; Juneau. Mr. Liebeck, who has kindly compared specimens with the Utah types in the Horn collection, finds certain small differences which seem to be constant and indicate that the Alaskan form may be distinct.

Laccobius ellipticus Lec. Anchorage.

Cercyon marinus Thoms. Dawson.

Cercyon fulvipennis Mann. Skagway; Anchorage; Juneau.

FAMILY SILPHIDÆ

Necrophorus pustulatus *var. melsheimeri* Kby. Anchorage.

Silpha lapponica Hbst. Anchorage.

Silpha trituberculata Kby. Dawson; White Horse.

Pelatines (*Pelates* Horn ||) *latus* Mann. Sitka.

Choleva spenciana Kby.? Dawson. Two females. *C. alsiosa* Horn is in my collection from Fort Yukon (Kusche). Although described from Alaska in 1885, this species does not appear in the Hamilton list of 1894.

Colon magnicolle Mann. McKinley Park. A single specimen.

Colon pusillum Horn. Dawson. Two examples.

Hydnobius substriatus Lec. Skagway; Dawson.

**Hydnobius latidens* Lec. McKinley Park.

Anisotoma punctatostriata Kby. Dawson.

**Anisotoma assimilis* Lec. Anchorage.

**Agathidium revolvens* Lec. McKinley Park; White Horse.

**Clambus gibbulus* Lec. Anchorage.

FAMILY SCYDMÆNIDÆ

Lophoderus biformis Mäkl. Sitka; Ketchikan.

**Euconnus longiceps* Fall, n. sp.

Of large size (2 mm.) and rather narrow form, polished, impunctate, except for the minute punctures bearing the elytral hairs; brown, with the sutural edge nigrescent basally. Pubescence yellowish-brown, stiff and posteriorly directed on the occiput, similarly rather densely bristling at sides of prothorax; longer, finer, sparser and more reclinate on the elytra. Head much longer than wide, scarcely more than half as wide as the prothorax; sides of the occiput nearly straight and convergent for about four times the length of the small anterior eyes, to the strongly constricted neck, the latter about one-half the width of the head. Antennæ fully half the length of the body, without definitely limited club; second joint subcylindrical, slightly wider apically, twice as long as wide and nearly equal to the two following; three to six equal in width to the second and mutually nearly similar, suboblong, slightly longer than wide; seven to ten gradually wider and becoming a little transverse; eleven not quite as long as the two preceding. Prothorax distinctly longer than wide, sides parallel and arcuate before the posterior lateral constriction, subbasal transverse impression feebly trifoveate at middle and at hind angles, the latter with short carina. Elytra elongate oval, widest at middle, about two-thirds longer than wide and two-thirds wider than the prothorax; basal impression strong, subhumeral and sutural impressions obsolete. Femora moderately clavate; tibiæ wider medially, tarsi rather short and stout, basal joint subequal to the two following; apical a little longer than the basal; hind coxæ subcontiguous. Length, 1.95 mm.; width, .75 mm.

Ketchikan, Alaska, August 1, 1924, a single example sifted from damp moss.

Although possessing the essential characters of *Euconnus*, this species does not fit very well in any of the groups of the genus established by Casey. The very elongate head should make it easily recognizable among all known species of the genus. The sex of the single individual is uncertain, but the slightly broader basal joint of the protarsus may indicate the male.

**Scydmaenus adjutor* Fall, n. sp.

Elongate, suboval, only moderately ventricose; black, antennæ and legs brownish ferruginous; surface polished, head and prothorax impunctate, elytra with fine sparse punctulation; pubescence fine, sparse and reclinate or inclined. Head wider than long; eyes rather prominent, subbasal, the tempora very short and rapidly convergent; excavation of the vertex broad and shallow, vaguely biimpressed at bottom. Antennæ gradually incrassate, the eighth joint slightly, the ninth and tenth more distinctly, transverse; joints proportionately nearly as in *pacificus*. Prothorax a little longer than wide, sides moderately rounded in apical half, constricted at basal third; surface coarsely biimpressed laterally within the constriction and with a

short subbasal dorsal transverse impression. Elytra oval, widest near middle, three-fourths wider than thorax, one-half longer than wide; humeri obtusely rounded but evident; basal impressions nearly as in *pacificus*; legs as in *pacificus*. Length (head depressed), 1.2 to 1.3 mm.; width, .55 mm.

Described from two examples taken by the writer at Skagway, Alaska, July 27, 1924.

This species is a *Brachycephsis*, closely allied to *subpunctatus* and *pacificus*. It is very considerably smaller than either of the two mentioned, and differs from *pacificus* also in its black color (rufotestaceous in *pacificus*), rather more slender antennæ and slightly less ventricose elytra, which are widest more nearly at the middle. There is no apparent indication of sex in either of the two specimens.

Eutheia (Veraphis) scitula Mäkl. Scow Bay.

FAMILY ORTHOPERIDÆ

**Orthoperus scutellaris* Lec. Dawson; Sheep Creek; Yukon River (below Eagle). This species was wrongly transferred to *Sphærius* in the Leng List.

**Sacium lugubre* Lec. White Horse; Dawson; Fort Yukon.

FAMILY STAPHYLINIDÆ

Olisthærus megacephalus Zett. McKinley Park; Tanana River.

Megarthrus sinuatocollis Lac. Dawson.

Anthobium pothos Mann. Seward.

**Pycnoglypta lurida* Gyll. Anchorage.

Omalium foraminosum Mäkl. Dawson; Sheep Creek, Yukon River (below Eagle).

Omalium sp. Fairbanks.

Phlæonomus lapponicus Zett. Dawson; Anchorage; Fort Yukon (Kusche).

**Arpedium brunnescens* Sahlbg. Skagway; McKinley Park.

Acidota quadrata Zett. Skagway; McKinley Park.

Microædus laticollis Mann. Sitka.

Microædus austinianus Lec. Seward; Ketchikan.

Pelecomalium testaceum Mann. Skagway; Seward.

Geodromicus ovipennis Lec. Dawson. One example. This specimen agrees well with LeConte's type and cannot safely be referred to the European *plagiatum* (see Leng list), a specimen of which is before me. This latter has the elytra distinctly longer, with less diverging sides.

Ancyrophorus biimpessus Mäkl. McKinley Park.

**Trogophlœus alaskanus* Fall, n. sp.

Moderately slender, depressed, sublinear; black, legs black, the tibiæ internally, tarsi, mouth and base of palpi testaceous; antennæ piceous. Upper surface densely, finely punctate and feebly shining; abdomen, especially toward apex, more sparsely punctate and shining; pubescence very short and fine. Head quadrate, hind angles subrectangularly rounded; tempora slightly longer than the eyes, straight and parallel; vertex broadly roundly impressed. Eyes small, just perceptibly more prominent than the tempora, attaining the middle of the elytra. Antennæ moderately slender, gradually incrassate, the outer three joints forming a feebly differentiated club; joints four to eight about as long as wide, fourth smallest, eighth just perceptibly transverse. Prothorax equal in width to the head, a little wider than long, front angles nearly right and scarcely rounded, sides broadly arcuate and feebly convergent from just behind the apex; hind angles obtuse and rounded as viewed from above; disc with an oblong impression occupying the apical two-thirds of the length and the middle one-third of the width. Elytra distinctly wider than thorax, subequal in length to the head and thorax combined and much longer than wide. Abdomen slightly narrower at base than the elytra and slightly widened behind, border rather thick but shallow; ventral segments four to six impressed at middle, the fourth very feebly, the sixth broadly and deeply, the apex of the latter behind the impression briefly carinate, the margin broadly angulate medially. Legs moderately slender. Length, 1.6 mm.; width, .45 mm.

McKinley Park Station, Alaska, July 9, 1924; a single example taken by the writer.

This little species belongs to Group IV of Casey's Revision, and by the table would be associated with *indigenus*, from which it differs in a number of details, notably the longer elytra, dark legs, and ventral impressions.

**Trogophlœus teres* Fall, n. sp.

Closely related to the preceding species, but apparently distinct by the following differences: Size larger (2.3 mm.), elytra piceotestaceous, legs entirely pale; prothorax slightly wider than the head and relatively more transverse, the impression much larger, extending almost throughout the length, at the base involving almost the entire discal width, but becoming narrower and more vague anteriorly. Elytra even slightly longer than in *alaskanus*, abdomen more distinctly narrowed basally and more strongly punctured, less shining. Ventral segments not impressed.

A single specimen only, taken at McKinley Park Station on the same date as the preceding, but whether the two specimens were taken together I do not recall. There is perhaps a possi-

bility that they are sexes of one species, the difference in color being due to immaturity. This point can only be settled by further collecting. So far as I know these are the first species of the genus to be recorded from Alaska.

Aploderus sp. Female. Ketchikan. This seems much like *princeps*, but without the male definite determination is difficult.

Oxytelus invenustus Csy. ? One female. White Horse. A male example taken at Edmonton, Alberta, on this trip, and others in my collection from Banff, Alberta, and Beulah, New Mexico, appear to possess precisely the male sexual characters described by Casey from his Maryland type.

**Oxytelus suspectus* Csy. Skagway.

Platystethus americanus Er. Dawson.

Bledius turgidus Csy. Yukon River, fifty miles below Selkirk, Y. T.

Bledius sp. Dawson.

Bledius albonotatus Mäkl. ? Skagway.

Stenus bipunctatus Er. Dawson.

**Stenus austini* Csy. McKinley Park; Anchorage. I have also a specimen of this species taken by Wickham at Fort Wrangel; this is, I suppose, one of the two undetermined species mentioned by Wickham in his list.

Stenus pterobrachys G. and H. Juneau; Sitka; Ketchikan.

**Stenus stygicus* Say. Eagle.

**Stenus mammops* Csy. Anchorage.

Stenus cariniceps Mäkl. Juneau; Ketchikan. Ten other as yet unidentified species of *Stenus* were taken in Alaska. I had hoped to be able before this to compare these with Casey's types, but Mr. Barber writes me that the latter are not yet available for study.

**Lathrobium sollicitum* Fall, n. sp.

Form nearly parallel; black, antennæ piceous, base scarcely paler; legs rufopiceous, the tibiæ paler. Head, thorax and elytra polished, abdomen dull from the density of the minute punctulation. Antennæ rather slender, not quite reaching the thoracic base, intermediate joints all feebly obconic and a little longer than wide. Head behind the clypeus subquadrate, sides parallel, hind angles broadly rounded, base truncate at middle, punctures numerous but well separated, median line smoother; eyes at about twice their length from the

base; gular sutures parallel and widely separated; neck one-third the width of the head. Prothorax slightly wider than the head, about one-fourth longer than wide, sides parallel, feebly arcuate, front angles rather abruptly, hind angles more broadly, rounded; punctuation strong, not very close, median line narrowly but indefinitely smooth. Elytra equal in length and in maximum width to the thorax, sides very feebly but perceptibly divergent from base to apex, punctures similar to those of the thorax but rather sparser. Abdomen at base equal in width to the elytra, very slightly widened to apical third, very finely, densely punctulate and dull, both above and beneath. Legs moderately stout, the front thighs thickened but not sinuate beneath. Length, 5.2 to 5.4 mm.; maximum width, .9 mm.

Described from two males taken by the writer at Anchorage, Alaska, July 13-14, 1924.

In these the sixth dorsal abdominal segment is strongly acuminate at apex, the sixth ventral with a very small apical emargination, triangular in form with its apex narrowly rounded.

By its narrow neck, this species should be referred to the group *Tartopeus*, but does not resemble at all any species therein defined by Casey, unless it be his *lacustris*, which is apparently quite different from the other species of his table, and is seemingly remindful of the more typical *Lathrobii*. In *lacustris* the elytra are said to be shorter than the thorax, and the sexual characters of the abdomen are not like those in *sollicitum*.

Lathrobium fnitimum Lec. Scow Bay.

Lathrobium subseriatum Lec. Kasaan.

****Lathrobium sewardi* Fall, n. sp.**

Form moderately slender and convex. Head, thorax, and basal half of elytra brownish piceous or castaneous, apical half of elytra brighter rufous; abdomen piceous, antennæ and legs reddish-brown. Head subquadrate, sides straight and parallel, hind angles broadly rounded, base truncate; eyes small, at twice their length from the base; surface densely punctate at sides and base, less closely so medially. Antennæ attaining base of prothorax, median joints obconical and about one-half longer than wide. Prothorax slightly narrower than the head, one-third longer than wide, widest at apex, sides nearly straight and convergent to base; all angles broadly rounded; punctuation close and confused, median impunctate line subelevated. Elytra as wide as the head, subequal in length to the prothorax, and one-sixth or one-seventh longer than wide; sides parallel, surface coarsely and closely sublinearly punctate. Abdomen very finely not closely punctate.

Male. Ventral segments two to six with median impressions in-

creasing in size and depth, margin of fifth segment broadly sinuate, apex of sixth with suboval or elliptical emargination, its limiting angles sharply subrectangular, the impression anteriorly with a dense patch of short black spicules, narrowly divided along the middle. Length, 5.8 to 7 mm.; width, .9 to 1 mm.

Described from two males and four females taken by the writer at Seward, Alaska, July 18-20, 1924.

This species belongs to Casey's genus *Lobrathium* and is very closely related to *tacomæ* Csy., differing most notably in color and the less slender antennæ; the secondary sexual characters are nearly identical.

Baptolinus macrocephalus Nordm. Ketchikan; Seward; Kasaan; Skagway.

Philonthus septentrionis Fall, n. sp.

Deep black throughout, polished, elytra with slightly green bronze luster. Head quadrate oval in male, a little more oval in the female; eyes distant their own length from the base of the head. Antennæ barely as long as the head and thorax, entirely black; joints eight to ten as wide or slightly wider than long, the preceding joints slightly more elongate. Prothorax quadrate oval, one-third to two-fifths wider than the head, evidently narrowed in front; dorsal punctures fine, three in number, equidistant. Elytra a little wider than the thorax, nearly square or very slightly transverse; punctuation rather fine and sparse, not distinctly muricate. Abdomen sparsely a little more strongly punctate than the elytra, beneath still more coarsely so.

Male. Anterior tarsi moderately strongly dilated; last ventral with a triangular emargination a little wider than deep, and surrounded by an acutely angulate gutter. Length, 6 to 7 mm.; width, 1.4 to 1.6 mm.

Described from one male and two female examples collected by the writer at Dawson, Yukon Territory, June 24, 1924. This species is of about the size and aspect of *umbratilis*, and may be placed near it. In the latter the tempora are shorter, the thorax less narrowed in front and with coarser serial punctures, the abdomen more finely punctured beneath, and the legs brownish-red.

Philonthus basalis Horn, var. Dawson. Three examples differing from the typical form by having the elytra varied with fuscous.

Philonthus fraternus Fall, n. sp.

Moderately elongate, deep black, shining, tarsi alone brownish.

Antennæ not quite as long as the head and thorax, outer joints nearly square. Head subquadrate, three-fifths as wide as the thorax, the latter as long as wide, distinctly narrowed in front; dorsal punctures four, equidistant. Elytra nearly square, a little wider than the thorax, not at all æneous, punctuation rather sparse. Abdomen a little more finely and equally sparsely punctate.

Male. Front tarsi very broadly dilated, last ventral triangularly emarginate and with a rather deep longitudinal impression tapering forward and extending the entire length of the segment; penultimate segment very broadly angulato-sinuate. Length, 6 mm.; width, 1.4 mm.

Dawson, Yukon Territory, June 24, 1924, a single male specimen taken by the writer.

This species can only be compared with *quadricollis* and *theveneti*, because of the last ventral of the male being longitudinally impressed throughout its length. In both these latter species the sides of the thorax are parallel or very nearly so, while in the present one they are distinctly convergent. *Quadricollis* has the legs and antennæ brown or paler and *theveneti* is said to have the elytra not wider than the thorax.

**Philonthus* sp. Ketchikan. One female in the vicinity of *crotchi*, which it may be.

**Philonthus distans* Horn. Anchorage.

Philonthus siegwaldi Mann. Skagway; Juneau; Port Althorp.

Philonthus nigritulus Grav. Skagway; Dawson.

**Philonthus decipiens* Horn. Wrangel.

**Philonthus sordidus* Grav. Anchorage.

Creophilus maxillosus L. Juneau.

Heterothrops sp. Near *carbonatus*, but probably distinct; Dawson, one female.

Quedius lævigatus Gyll. Seward.

Quedius molochinus Grav. Seward; Skagway; Juneau.

Quedius brunneipennis Mann. Anchorage; McKinley Park.

Quedius hyperboreus Er.? Anchorage.

Quedius sublimbatus Mäkl. Skagway.

Quedius sp. Childs Glacier.

Quedius peregrinus Grav. Juneau.

Quedius sp. near *ænescens* Mäkl. Fairbanks.

Tachinus nigricornis Mann. Wrangel.

Tachinus instabilis Mäkl. Anchorage; Juneau.

- Tachinus circumcinctus* Mäkl. Dawson; Tanana River; Fort Yukon (Kusche).
- **Tachyporus jocosus* Say. Anchorage.
Tachyporus chrysomelinus Er. Dawson.
Boletobius cincticollis Say.? Anchorage; McKinley Park.
Mycetoporus humidus Say. Eagle.
Mycetoporus nigrans Mäkl. McKinley Park, one example which fits Mäeklin's description very perfectly. His type was said to be from the interior of the Kenai Peninsula.
- **Gymnusa variegata* Kies. Fairbanks, one example, differing slightly from a European specimen in my collection.
- **Myllæna* sp. Scow Bay; Ketchikan. No species of this genus has hitherto been recorded from Alaska.
Placusa complanata Er. Dawson.
Gyrophæna sp. Anchorage. This is perhaps *geniculata* Mäkl.
- **Atheta divisa* Märk. Anchorage; McKinley Park.
Atheta aquatica Thom.? Dawson.
Atheta graminicola Grav. Skagway; Fairbanks; White Horse.
Atheta picipennis Mann. Anchorage.
Acrotona fungi Grav. Skagway; Anchorage; Juneau.
- **Gnypeta* sp. Fairbanks. First record of the genus from Alaska.
Gnypeta sp. Dawson.
Aleochara bipustulata L. Anchorage; Fairbanks; Dawson.
Aleochara pleuralis Csy. Dawson.
Dasyglossa prospera Er. Tanana River. The greater number of the Aleocharinæ taken are as yet undetermined.

FAMILY PSELAPHIDÆ

- Sonoma parviceps* Mäkl. Ketchikan.
Trimium clavicorne Mäkl. Ketchikan, one pair.
- **Richenbachia binodifer* Csy. Ketchikan. Scow Bay.
 **Rybaxis brendeli* Lec. var. Anchorage.
 **Tychus cognatus* Lec. Seward; Ketchikan.

FAMILY PTILIIDÆ

- Ptenidium pullum* Mäkl. Scow Bay.
Acratichis (Trichopteryx || Kby.) sitkensis Mots. Sitka.
Acratrichis laticollis Mäkl. Seward.
Acratrichis sp. Anchorage.

Acratrichis sp. McKinley Park.

Acratrichis three species. Dawson.

FAMILY HISTERIDÆ

Hister (Psilocelis) subopaca Lec. var., or very closely allied species. Dawson, one example; White Horse (Kusche).

FAMILY LAMPYRIDÆ

**Podabrus comes conspiratus* Fall. n. var.

This name is proposed for a form of which a single example was taken at Skagway. It may or may not be a form of the Californian *comes*, the amount of variability of which within specific limits is not definitely known. The present specimen differs from all my *comes* (so labelled) by the slightly more transverse thorax, which is a little less rounded on the sides and without visible sinuation at the hind angles, which are not perceptibly prominent. The thorax also lacks the median eroded line and has the front margin blackish at middle. It agrees otherwise in color with the typical *comes* except that the posterior ventral segments are not margined with yellow, the terminal segment being merely feebly paler at tip. Length, 14 mm.

Skagway, Alaska, June 18-20, 1924; a single female.

**Podabrus tetragonoderus* Fall. n. sp.

Moderately elongate, black, clypeus at sides, sides of prothorax and basal three joints of antennæ beneath, yellow; elytra with a grayish aspect due to the rather plentiful ashy pubescence. Antennæ longer than half the body; second joint a little narrower but subequal in length to the third (♂), or a little shorter than the third (♀); third joint subtriangular, a little longer than wide, distinctly wider than the fourth in the male, less noticeably so in the female; following joints nearly parallel-sided, becoming gradually more slender, the fourth a little more than twice, the outer joints fully three times, as long as wide. Head (♂) just perceptibly narrower than the thorax; eyes moderate; the eyes less prominent and the head narrower in the female; clypeus and front as far back as a line joining the posterior part of the eyes nearly smooth; occiput closely, rugosely punctate but shining. Prothorax quadrate, a little wider than long, not or barely perceptibly wider near the front angles; sides nearly parallel and straight; a very faint sinuation before

the sharply defined hind angles, front angles rounded, posterior discal elevations rather strong and situated within the broad dorsal black stripe, which is dilated behind the middle; surface distinctly but not deeply punctate throughout, more finely so in the yellow margins and between the discal elevations; median line finely eroded in posterior two-thirds. Elytra finely scabrous punctate. Legs black, front tibiæ in the single female a little dilute in color. Claws (♂) finely cleft, the inner portion approximate to, and nearly as long as, the outer on the front and middle feet, much shorter but still finely acute on the hind feet. In the female the inner portion of the claw is much shorter than the outer on all feet, and the claws might perhaps better be described as having a basal dilatation with the free angle sharply acuminate. Length, 7 to 9.5 mm.

Described from three males and one female, taken by the writer at Skagway, Alaska, June 18-20, 1924.

This species will not enter any of LeConte's groups as defined though coming nearest to Group C. There are other species in my collection with unguual characters similar to the present one and I hope shortly to publish already prepared descriptions of these and other new species of the genus.

***Podabrus fissilis** Fall, n. sp.

Black, feebly shining; muzzle and two or three basal joints of antennæ beneath rufotestaceous. Antennæ slender; second joint but little more than half the third in the male, a little shorter than the third in the female. Head conspicuously wider than the thorax (♂), or subequal in width to it (♀); surface alutaceous and opaque, front sparsely punctate. Prothorax small, nearly as long as wide in the male, front angles obliquely subtruncate, sides a little convergent thence to the somewhat prominent hind angles. There is a small rufescent marginal spot just behind the front angles in the male which is entirely lacking in the female. Prothorax finely punctate and alutaceous, sculpture as usual. Elytra very finely scabrous; tarsal claws all finely cleft in the male, the inner part shorter on the posterior four feet; all claws toothed at base in the female. Length, 6 to 7 mm.

Described from a single pair (type male) collected at Emerald Lake, British Columbia, by Dr. A. Fenyès. This is

the first nearly entirely black species to be described having tarsal claws all finely cleft in the male.

Since writing the above description I have taken a single male of this species at Skagway, Alaska.

Podabrus piniphilus Esch. Skagway; Seward. The Seward example (male) may be different.

**Podabrus extremus* Lec. Skagway.

**Podabrus lævicollis* Kby. Anchorage.

FAMILY CANTHARIDÆ (TELEPHORIDÆ)

Cantharis (Telephorus) mandibularis Kby. (= *nigritulus* Lec.). Eagle; Skagway; McKinley Park; Dawson; White Horse. This is a common species in Alaska and ranges across the northern part of the continent to New Hampshire. *Mandibularis* has long been erroneously placed as a synonym of *fraxini* in our literature, but the above synonymy is definitely established by comparison with one of Kirby's cotypes.

**Silis difficilis* Lec. Skagway.

FAMILY MELYRIDÆ

Dasytes hudsonicus Lec. Skagway; White Horse; Dawson; Eagle; Fort Yukon; commonly beaten from spruce.

FAMILY CLERIDÆ

Thanasimus undulatus Say. Dawson.

Trichodes ornatus Say. Dawson.

FAMILY CORYNETIDÆ

**Necrobia violacea* L. Dawson; Fort Yukon (Kusche).

FAMILY MORDELLIDÆ

**Anaspis atra* Lec.? Dawson; Eagle.

Anaspis sericea Mann. Skagway; Sitka; Seward.

FAMILY PYTHIDÆ

Priognatha monilicornis Rand. Skagway.

FAMILY ANTHICIDÆ

Anthicus melancholicus Laf. Dawson.

FAMILY ELATERIDÆ

Athous ferruginosus Esch. Skagway; Sitka; Ketchikan. As Dr. Van Dyke has pointed out in his recently published Katmai List, this is but a slight variety of *rufiventris* Esch., which is the older name.

(To be continued)

TWO NEW SPECIES OF THRIPS FROM
CALIFORNIA

BY ARTHUR C. MASON

*Bureau of Entomology, U. S. Department
of Agriculture*¹**Hoplandrothrips sycamorensis** Mason, new species

Female. Length, 2.11 mm. Color uniformly dark brown, with a very little scattered red hypodermal pigmentation; tarsi, fore tibiae, and bases of antennal segments III to VI light brown.

Head nearly 1.5 times as long as broad, widest immediately behind the eyes and converging slightly posteriorly; frons overhanging about half of the first antennal segment. Cheeks nearly straight and somewhat roughened with small warts, and having several very short blunt spines. Postocular bristles short, dilated, and truncate at tip. Ocelli small, circular and equidistant, placed between the eyes; anterior one near the end of the vertex and the posterior pair about opposite the center of the eyes. Eyes small, occupying little more than one-fourth of the length of the head. Mouth-cone long, almost reaching the mesothorax and running to a sharp point. Antenna eight-segmented and one and one-fourth times the length of the head. First two segments concolorous with the head, III light yellow with a little brown shading near the tip, IV to VI yellow at base, shading into brown, VII and VIII light brown, VII having a very little yellow at base. Segment I a short truncated cone, II cup-shaped with a rather pronounced pedicle, III wedge-shaped with rounded apex, IV and V ovate with narrowed pedicle, IV being considerably wider than V, VI fusiform, VII truncate ovate, VIII conical with a blunt tip.

Prothorax little more than half as long as the head and about twice as wide as long, sides diverging rapidly posteriorly; one rather prominent but short spine on the anterior margin and a row of five spines on the posterior margin, the first of them being especially heavy. All spines dilated and truncate at tip.

Pterothorax slightly wider than the prothorax and narrowed somewhat posteriorly. Wings rather short and colorless, except for a little yellow shading at the extreme base; three blunt spines at base of fore wings, five to nine (usually eight) interlocated hairs on posterior margin. Legs rather long, the fore femora being heavy and thickened; fore tibiae and extreme tips of others light yellow. Fore tarsi armed with a short, stout tooth near base.

Abdomen large, tapering gradually after the fourth segment and rounded abruptly from the seventh; provided with numerous blunt

¹ Published with the permission of the Secretary of Agriculture.

colorless bristles, the posterior ones and those at the end of the tube being nearly as long as the tube. Tube half as long as the head and tapering gradually.

Measurements: Head, length 0.32 mm., width 0.23 mm.; prothorax, length 0.18 mm., width 0.37 mm.; pterothorax, width 0.38 mm.; abdomen, greatest width 0.408 mm.; tube, length 0.15 mm., width at base 0.073 mm., width at apex 0.043 mm.; antenna, total length 0.41 mm.

Segment	1	2	3	4	5	6	7	8
Length	34	54	67	72	59	49	46	28 microns
Width	43	35	38	39	32	29	25	16 microns

Male slightly smaller and with more red pigmentation; otherwise similar in appearance. Length, 2.03 mm. As with the females, the fore tarsi are provided with a stout tooth and in addition they have a smaller tooth on the inside of the fore tibiae near the base and also a very small one opposite this on the femur.

The larvæ are light yellow to colorless, and live under the bark of the host. Winter is passed in the egg stage, under the bark.

Described from five females and one male taken under bark or on the leaves of sycamore trees at Springville, California. Type in author's collection.

This species is close to *H. xanthopus* Hood, but differs in many characters, including larger size, darker color, absence of most of the red pigmentation, shorter postocular and prothoracic bristles, etc.

Zygothrips californicus Mason, new species

Length, 1.35 mm. Color brown to dark brown or fuscous; legs concolorous with the body; antennal segment III yellow, IV to VIII light brown.

Head slightly longer than broad, rounded in front and slightly diverging posteriorly, reaching the widest part about three-fourths of the way back, then narrowing slightly to the posterior end. Cheeks smooth. Postocular bristles very small and blunt. Eyes rather small, occupying about one-fourth of the length of the head and one-third of its width. Ocelli rather large, the posterior pair being opposite the anterior third of the eyes and the anterior one being directed forward. Mouth-cone contracted, reaching about two-thirds across the pronotum, bluntly rounded. Antenna eight-segmented, one and one-half times the length of the head; bristles short but sense-cones rather conspicuous. First two segments concolorous with the head; III light yellow; IV to VI brownish-yellow to light brown; VII and VIII light brown. Segment I a short truncated cone; II ovate; III clavate with a broad base; IV ovate, tapering abruptly to a narrow base; V barrel-shaped; VI cylindrical, with nar-

rowed base; VII oblong and slightly tapered at tip; VIII conical, tapering gradually to a rather sharp point.

Prothorax about three-fourths as long as the head and nearly twice as wide as long, the sides gently diverging and broadly rounded at the posterior end; prothoracic bristles short and blunt.

Pterothorax slightly wider than the prothorax, sides nearly straight and converging slightly posteriorly. Wings large and clear, with long membrane extending to or slightly beyond the end of the tube. Fore pair somewhat constricted in the middle and with nine to eleven interlocated hairs on the posterior border. Legs long, femora enlarged, the fore femora especially being much thickened; fore tarsi unarmed.

Abdomen rather heavy, width about four-ninths of the length, and tapering rather abruptly from the fifth segment to the tube. Tube less than half the length of the head; terminal bristles little more than half the length of the tube.

Measurements: Head, length 0.229 mm., width 0.209 mm.; prothorax, length 0.152 mm., width 0.276 mm.; pterothorax, width 0.304 mm.; abdomen, greatest width 0.342 mm.; tube, length 0.098 mm., width at base 0.057 mm., width at apex 0.033 mm.; antenna, total length 0.328 mm.

Segment	1	2	3	4	5	6	7	8
Length	25	41	49	56	49	41	41	21 microns
Width	32	29	29	37	29	25	18	10 microns

Described from a single female taken on an orange tree at Orange Cove, California. Type in author's collection.

Close to *Z. harti* Hood, but differing in the following characters: Larger size, antennal segments IV to VIII lighter colored, prothoracic bristles blunt, clear fore wings and with 9 to 11 interlocated hairs, contracted mouth cone, and tube less than half the length of the head.

NOTES ON PARASITES AT SATICOY, CALIFORNIA, DURING THE YEAR 1925

BY STANLEY E. FLANDERS

Saticoy, California

On February 28 a braconid of the genus *Monogonogastra* was reared from a walnut twig which contained the dead larvæ of *Polycaon confertus* Lec. During April numbers of these dusky-winged, red-bodied braconids were observed about the walnut leaves. One individual was seen December 4.

Through June and July a number of the metallic-blue ichneu-

mon-flies *Amblyteles cæruleus* (Cr.), and the yellow and black *A. zebratus* (Cr.) were reared from the pupæ of *Sabulodes forficaria* Guen.

Hovering about the blossoms of a large eucalyptus tree on April 2 were many of the syrphid parasites, *Homotropus decoratus* (Cr.).

During October a black chalcid, *Zatropis tortricidis* Crawford, was reared from the lima-bean pod-borer, *Etiella zinckenella* Treit. It pupates free in the pod, and is in the pupal stage seventeen days.

On June 11 the gray, oval larvæ of a species of *Euplectrus* were observed in a cluster on the dorsum of a caterpillar of *Sabulodes forficaria*. Later, several of these parasitized caterpillars were found.

From the pupæ of the lichen caterpillar, *Illice nexa* Boisd., a number of specimens of the tachinid, *Pseudochæta argentifrons* Coq., were reared in May.

On June 20 three females of *Epiurus* sp. (possibly *Pimpla albipes* Cr. described from Mexico) were reared from *Archips argyrospila* Walk. Their silken cocoons were spun on walnut about the remains of the caterpillar.

Six small chalcids of the genus *Elachertus* were reared in June from a caterpillar of *Archips argyrospila*.

On April 29 a walnut aphid, *Chromaphis juglandicola* Kalt., was found parasitized by *Aphelinus perpallidus* Gahan. It is only occasionally that such a parasitized aphid is found.

While collecting codling moth larvæ from burlap bands on walnut trees during March, a light brown ichneumon-fly, *Exetastes* n. sp., was taken from a codling moth cocoon.

Between May 1 and July 21 twenty-four tachinids of the genus *Lixophaga* were reared from three hundred overwintering codling moth larvæ. The maggot pupates within twelve hours after leaving the caterpillar. The length of the pupal stage was nineteen and twenty days.

The identification of the parasites noted were kindly made by the specialists, R. A. Cushman, A. B. Gahan, S. A. Rohwer, J. M. Aldrich, and P. H. Timberlake.

SEX-LIMITED POLYCHROMATISM IN
LASIOPHTICUS PYRASTRI (LINN.)

BY E. L. KESSEL

Early last June I collected a large population of the common syrphid fly *Lasiophticus pyrastris* (Linn.) on the University Farm at Davis, California. This series, which was taken by sweeping so as to be representative of the fauna, numbered 277 individuals. When a classification was made, there were found to be 235 specimens of the normal bicolored phase, of which number 98 were females and 137 were males. This is a decidedly unbalanced sex ratio. The remaining 42 members of the population were of the melanic variety, described by Curtis as *unicolor*. It was observed that all of these melanic flies were females, and it is interesting to note that when their number is added to the 98 females of the bicolored type there appears the well-balanced sex ratio for the whole population of 140 females and 137 males.

The above data lead to the conclusion that the melanic form *unicolor* appears only in the female sex. Moreover, Curtis, in his description of this variety, gives the impression that he had but one specimen, and his colored illustration is definitely that of a female. Still more evidence was obtained from an examination of the material in the collection of the California Academy of Sciences. This series contains thirteen specimens of *unicolor* collected in various parts of California, and all of these are females. The evidence, therefore, appears to be conclusive.

An intermediate form, although comparatively rare, also occurs. I took three specimens of this variety in Davis last June, but these were obtained by selective collecting. They are characterized by having the yellow markings persisting on the second abdominal segment, although they are much reduced in size. The California Academy of Sciences has one such individual collected by Dr. E. C. Van Dyke on Mount Ranier, Washington, in 1920. Two other specimens are cited by Cole and Lovett in their "Diptera of Oregon." All six of these intermediates are also females. We may, therefore, assume that this variety, as well as the completely melanic one, is sex-limited to the female. The species, therefore, appears to be an example of sex-limited polychromatism, in which there is only one type of male but three types of females.

THE PAN-PACIFIC ENTOMOLOGIST

Published quarterly by the Pacific Coast Entomological Society
in co-operation with the California Academy of Sciences

E. P. VAN DUZEE, Editor F. E. BLAISDELL, M. D., Treasurer

Through an oversight the legend was omitted from the plate of *Hypolimnas bolina* L. in the October number of this journal. A gummed slip to be attached to this plate is enclosed.

On March 23 the Pacific Coast Entomological Society gave a complimentary dinner to Dr. and Mrs. William Barnes of Decatur, Illinois, and Dr. and Mrs. R. J. Tillyard of Nelson, New Zealand. Dr. Barnes gave the society an interesting account of his work on the North American Lepidoptera and Dr. Tillyard on his work at the Cawthorn Institute of Scientific Research at Nelson. Dr. Barnes spent several days at the Academy of Sciences, and his suggestions added much of value to the Academy collection of Lepidoptera.

The recent death of Mr. Richard H. Stretch at his Seattle home has removed one of the few remaining landmarks of early western entomology. About fifty years ago he amassed a very complete collection of the Bombycine moths of the world. This he presented to the University of California. So much of this collection as has escaped the ravages of pests has recently come into the care of the California Academy of Sciences as a permanent loan, where it will be accessible to those interested in the published work of Mr. Stretch.

Mr. E. Piazza, well known to our Lepidopterists as a successful collector of moths, died in New York, March 9, 1926, while on his way to England to visit his sister. Although an invalid for years he has been able to do much valuable work in the collection of moths for Scientific Institutions.

Through the generosity of friends we are able to give our subscribers sixteen extra pages in this number of the *Pan-Pacific Entomologist*. This enables us to include two larger papers that otherwise it would have been impossible to publish in full. The April number should follow this with little delay. The October number was mailed January 16, 1926.

Dr. F. E. Blaisdell writes us that it will be impossible for him to determine Coleoptera for his correspondents after September 1, 1926, and asks that we call the attention of coleopterists to this notice.

WARD'S NATURAL SCIENCE ESTABLISHMENT

College Avenue, Rochester, New York

We offer our best-made genuine Schmitt boxes, Exhibition cases, Cabinets, Insect pins, Riker mounts, Insects and Insect collections, and all other supplies essential for the pursuit of the study of entomology. Send for free supply catalogue, No. 41.

FOR SALE
PACIFIC COAST ENTOMOLOGICAL SOCIETY
PROCEEDINGS

Vol. I (1st to 80th meetings). 1901 to 1921. 187 pages. \$5.

Address: Dr. F. E. Blaisdell, 1520 Lake Street,
San Francisco, California

-
- From California Academy of Sciences, Proceedings, Series 4
- Williams, F. X., Butterflies and Hawk-Moths of the Galapagos Islands. 33 pages, colored plate. 1911. 50 cents.
- Hebard, Morgan, Dermaptera and Orthoptera (of the Galapagos Islands). 35 pages, Illust. 1920. \$1.
- Ferris, G. F., A Catalogue and Host List of the Anoplura. 84 pages. 1916. 50 cents.
- Essig, E. O., and Kuwana, S. I., Some Japanese Aphididæ. 77 pages, Illust. 1918. 45 cents.
- Aldrich, J. M., Kelp Flies of North America. 22 pages, Illust. 1918. 25 cents.
- Cole, F. R. and Lovett, A. L., New Oregon Diptera. 34 pages, 6 plates. 1919. 40 cents.
- Hanna, G. D. and others, Insects of the Pribilof Islands, Alaska. 42 pages, Illust. 1921. 50 cents.
- Cole, F. R. and Lovett, A. L., An Annotated List of the Diptera of Oregon. 147 pages, Illust. 1921. \$1.
- Curran, C. H., Revision of the Pipiza Group of the Syrphidæ from North America. 48 pages, Illust. 1921. 60 cents.

Address: California Academy of Sciences,
San Francisco, California, U. S. A.



THE PAN-PACIFIC ENTOMOLOGIST

Published by the
Pacific Coast Entomological Society
in co-operation with
The California Academy of Sciences

CONTENTS

FREEBORN, A NEW CHAOBORID GNAT	161
VAN DUZEE, A NEW SPECIES OF SCENOPINIDÆ	164
MACDOUGALL, SOME NEW SPECIES OF MACROSIPHUM FROM BRITISH COLUMBIA	165
DAVIS AND SMITH, NOTES ON THE GENUS STENOPELMATUS	174
FOX, SOME NEW SPECIES OF SIPHONAPTERA	182
BARNES AND BENJAMIN, ON THE PLACEMENT OF "PSEUDANARTA" FLAVI- DENS GRT.	187
BARNES AND BENJAMIN, ON THE DISTRIBUTION OF OMPHALOCERA OCCI- DENTALIS B. AND B.	188
TANNER, A NEW SPECIES OF PLASTOCERIDÆ IN EUTHYSANIUS	188
FALL, A LIST OF THE COLEOPTERA TAKEN IN ALASKA IN 1924	191
BEAMER, NOTES ON GRIBURIUS MONTEZUMA SUF.	209
ESSIG, A BUTTERFLY MIGRATION	211
FOX, A NEW NOMADA	212
DOANE, CONTROLLING MEALYBUGS ON ORNAMENTAL PLANTS	213
EDITORIAL	216

San Francisco, California

1926

THE PAN-PACIFIC ENTOMOLOGIST

Published quarterly in July, October, January and April by the Pacific Coast Entomological Society in co-operation with the California Academy of Sciences.

Annual subscription \$2.00 in advance for the United States and Canada; \$2.25 for foreign countries. Subscriptions should be sent to the treasurer, Dr. F. E. Blaisdell, 1520 Lake Street, San Francisco, California. Make checks payable to the "Pan-Pacific Entomologist."

Manuscripts for publication and communications regarding non-receipt of numbers, change of address, requests for sample copies, etc., should be addressed to the editor, Mr. E. P. Van Duzee, California Academy of Sciences, Golden Gate Park, San Francisco, California. Advertisements will be accepted for the back cover pages. For rates address the editor or treasurer.

Twenty-five copies of author's extras will be furnished free on request. Additional copies will be supplied at cost of publication if a request is received with the manuscript.

Subscribers failing to receive their numbers will please notify the editor at as early a date as possible.



PUBLICATION COMMITTEE, PAN-PACIFIC ENTOMOLOGIST

E. O. ESSIG, *Chairman*

G. F. FERRIS

R. A. DOANE

E. C. VAN DYKE

GRANT WALLACE

REGIONAL MEMBERS

W. W. HENDERSON, Logan, Utah

J. C. CHAMBERLIN, Riverside, California

E. P. VAN DUZEE, *Editor*

E. C. VAN DYKE, *Associate Editor*

F. E. BLAISDELL, SR., *Treasurer*

Published at the California Academy of Sciences, Golden Gate Park, San Francisco, California.

Entered as second-class matter, February 10, 1925, at the postoffice at